

# What is Suboxone?

The sublingual tablets called [Suboxone](#) include buprenorphine and naloxone. It includes naloxone HCL dihydrate, an opioid receptor antagonist, in a 4:1 ratio with buprenorphine HCl, a partial agonist at the mu-opioid receptor.

The drugs buprenorphine and naloxone are both present in this medication. It is employed to treat opioid addiction and dependence. The medication buprenorphine is a member of the mixed opioid agonist-antagonist pharmacological class. The withdrawal symptoms that come with discontinuing other opioids are lessened by [buprenorphine](#).

When administered, the opioid antagonist naloxone can result in very bad opioid withdrawal symptoms by blocking the effects of opioids.

When administered orally or dissolved under the tongue, it has little impact. In order to prevent drug abuse and misuse (injection of this medication), it is coupled with buprenorphine.



In a comprehensive programme to treat drug misuse, this combo medicine is employed (such as compliance monitoring, counseling, behavioral contract, lifestyle changes).

## Uses of Suboxone –

Before beginning to use buprenorphine/naloxone and each time you get a refill, read the medication guide that your pharmacist has provided. Ask your physician or pharmacist if you have any queries.

There are numerous brands and dosage options for this drug. Do not switch brands or dose forms without first consulting your doctor or pharmacist because various products may differ in their dosages of buprenorphine and naloxone.

Use this drug as prescribed by your doctor, typically once daily, during the treatment maintenance phase. Allow the drug to thoroughly dissolve by placing it under your tongue for 5 to 10 minutes.

If your doctor has given more than one tablet each day, you can either put them all under your tongue at once or two at a time. Never chew or swallow this drug. It won't function as well.

Your medical condition and treatment response will determine the dosage. Never use the medication in larger amounts or for longer than recommended. Also, never take it more regularly. When instructed, discontinue taking the medication properly.

Order Now/ 10% Discounts/ online order

<https://buyxanaxshop.com/product/suboxone-2mg/>

<https://buyxanaxshop.com/product/suboxone-8mg/>

To reap the greatest benefits from this drug, use it frequently. Use it at the same time every day to aid in memory.

If you use buprenorphine/naloxone soon after taking opioids like heroin, morphine, or methadone, you may experience withdrawal symptoms. For your treatment regimen, adhere to your doctor's directions.

You can get withdrawal symptoms if you abruptly stop taking this drug (such as restlessness, watering eyes, runny nose, nausea, sweating, muscle aches). Your doctor might gradually reduce your dose in order to help prevent withdrawal. If you have taken this medication for a long period or in large doses, withdrawal is

more likely to occur. If you experience withdrawal, tell your doctor or pharmacist straight away.

Buprenorphine/naloxone should not be injected or inhaled through the nose ("snort").



Due to the naloxone in this drug, doing so is risky and may result in severe withdrawal symptoms (see the Side Effects section), especially if you have previously used opioids like heroin, morphine, or methadone.

## Opioid Addiction:

It is advised to use Suboxone as a treatment for opioid addiction. By lessening the reliance and withdrawal brought on by the opioids, it aids in the reduction of opioid addiction.

## Side effects

There could be drowsiness, vertigo, constipation, or headaches. Inform your doctor or pharmacist as soon as possible if any of these side effects persist or get worse.

Consume dietary fiber, get adequate water, and exercise to avoid constipation. You might also require a laxative. Find out which kind of laxative is best for you by asking your pharmacist.

When getting out of a sitting or lying posture, take it gently to lessen the chance of feeling lightheaded and dizzy.

Keep in mind that your doctor ordered this medicine because, in his or her opinion, it will benefit you more than it will harm you. Many users of this medicine report no significant negative effects.

If this medicine is misused, injected, or used with other depressants, severe (potentially fatal) respiratory issues may develop (such as alcohol, benzodiazepines including diazepam, other opioids).

If you have any of these unlikely but dangerous side effects, contact your doctor immediately away: sleep apnea, mental/mood problems (such as agitation, disorientation, or hallucinations), stomach/abdominal pain, or indications that your adrenal glands aren't functioning properly (such as unusual tiredness, weight loss).

The adverse reaction by the Suboxone tablets include:

- Addiction, Abuse, and Misuse.
- Whole-Body: asthenia, chills, headache, back pain, abdominal pain, opioid withdrawal, hypersensitivity reactions, and neonatal opioid withdrawal syndrome.
- Cardiovascular: vasodilation.
- Digestive System: constipation, diarrhea, vomiting, and nausea.
- Nervous System: insomnia and CNS (Central Nervous System) depression.
- Respiratory System: rhinitis and respiratory depression.

**Order Now :** <https://buyxanaxshop.com/product-category/buy-suboxone-online/>